



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Is malnutrition associated with orthopaedic infections? A single-centre pilot evaluation

Guanziroli, N ; Hannouche, D ; Uçkay, I

DOI: <https://doi.org/10.1016/j.jhin.2018.11.006>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-158521>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Guanziroli, N; Hannouche, D; Uçkay, I (2019). Is malnutrition associated with orthopaedic infections? A single-centre pilot evaluation. *Journal of Hospital Infection*, 101(2):229-230.

DOI: <https://doi.org/10.1016/j.jhin.2018.11.006>

Accepted Manuscript

Is malnutrition associated with orthopaedic infections? A single-centre pilot evaluation

N. Guanziroli, D. Hannouche, I. Uçkay

PII: S0195-6701(18)30593-0

DOI: <https://doi.org/10.1016/j.jhin.2018.11.006>

Reference: YJHIN 5593


To appear in: *Journal of Hospital Infection*

Received Date: 21 August 2018

Accepted Date: 6 November 2018

Please cite this article as: Guanziroli N, Hannouche D, Uçkay I, Is malnutrition associated with orthopaedic infections? A single-centre pilot evaluation, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2018.11.006>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Is Malnutrition Associated with Orthopedic Infections: a Single-Center Pilot Evaluation

Nastassia Guanziroli¹, Didier Hannouche¹, Ilker Uçkay^{1,2}

¹Orthopedic Surgery Service, Geneva University Hospitals, Geneva, Switzerland

²Infectiology, Balgrist University Hospital, Zurich, Switzerland

Corresponding author

Ilker Uçkay, MD

Balgrist University Hospital

Forchstrasse 340, 8008 Zurich / Switzerland

E-mail: ilker.uckay@balgrist.ch

All authors waive financial support, grants, or royalties and have no financial interests that could lead to a conflict of interest. All authors state that they have read and approved the manuscript. It has not been published elsewhere nor is it under consideration for publication elsewhere. Parts of the manuscript have been presented as a poster at the 5th Oxford Bone Infection Conference, United Kingdom, 22-23 March 2018.

Keywords: *malnutrition; orthopedic surgery; association: infections*

Running title: Malnutrition and orthopedic infections

Sir,

Malnutrition is frequent among adult surgical patients [1-3]. Some authors suggest that malnutrition may be an independent risk factor for orthopedic infections. There are, however, few strong original data addressing this issue. We thus performed a prospective pilot quality evaluation for patients at risk of surgical site (SSI) or community-acquired orthopaedic infections. During February 2017-March 2017, we prospectively surveyed the nutritional status of 23 infected (nosocomial and community-acquired infections) and 18 uninfected patients, consecutively hospitalized in our orthopaedic Service. To estimate the presence of malnutrition, we used nine different clinical and laboratory parameters such as the Mini Nutritional Assessment Tool (MNA[®]) of the Nestlé Nutrition Institute [4], the body mass index, weight, history of weight loss in the last three months, presence of gastrointestinal diseases or diabetes mellitus, serum albumin levels, serum total protein and the serum pre-albumin levels at admission [1,2]. Additionally, we observed if the patient finished his/her meals during hospitalization. As a main result, we found no single difference between those with and without infections in any of these nine standard nutritional parameters (Table 1).

We cannot fully explain this complete absence of difference. Many colleagues think that there is a strong scientific link between malnutrition and SSI. The mechanism by which malnutrition would increase the risk for SSI equally remains obscure. In reality, available scientific literature is scant and national and international guidelines attribute a very low quality of evidence for this association. For example, the CDC guidelines of 1999 remain vague [5]. The recent global WHO guidelines [6] state that *"The panel suggests considering*

the administration of oral or enteral multiple nutrient-enhanced nutritional formulas for the purpose of preventing SSI in underweight patients [only] who undergo major surgical operations", taxing the strength of their own recommendations as "conditional" and the quality of evidence as "very low".

It might be that the value of the current diagnostic schema for malnutrition may lie essentially in identifying sicker patients. To cite a frequent example, most author groups use hypoalbuminemia with a cut-off of 3.5 g/dL [1,7,8] as the most important marker for malnutrition. However, although albumin may be a reliable serum marker of chronic nutritional status, it is not specific, in that inflammation or stress may also cause hypoalbuminemia without associated malnutrition [1]. Likewise, malnutrition is a confusing and complex term. It includes phenomena related to starvation, includes patients who are overweight as well as those with low nutritional intake. While obesity is a clear risk factor for SSI [1], other aspects of malnutrition might not be.

Our study has three important limitations. Without doubt, the sample size would be too small compared to a randomized trial or a case-control study. Indeed, our study cannot address interventional issues and it is unlikely to be valuable as a platform from which to launch a full study of the topic. In contrast, this study was a prospective pilot evaluation in preparation of a possible interventional trial by searching the best parameters to influence in such a possible future trial. Secondly, in our evaluation, even if we compared nine different parameters, we omitted other literature markers such as total lymphocyte count, serum zinc levels, iron/transferrin, pyridoxin and vitamin-D levels, lymphocyte/monocyte ratios, the measurement of healing time, or anthropometric measurements such as calf and arm

muscle circumference or triceps skinfolds (which all are not routinely assessed on admission). These are all markers of both, malnutrition and SSI, alike. In future trials, we would also need to consider not only gross protein and carbohydrate intake but also (and perhaps especially) micronutrients [1,8]; and perhaps to concentrate on nutritional status independently of confounders such as diabetes and high alcohol intake, gastrointestinal disease and immune-suppression. Accepting that these are important, any proposed trial to determine whether preoperative nutritional optimization might be more feasible without their inclusion. Finally, a literature bias in favor of positive associations is not entirely excluded. Negative results might be less prone for publication. Our pilot study is such a negative evaluation.

Ethical approval

Due to its small sample size, the non-invasiveness, and the use of already sampled parameters at admission, our quality evaluation was waived from formal written consent.

Conflict of interest statement

All other authors declare no conflicts of interest.

Funding sources

None.

References

1. Ellsworth B, Kamath AF. Malnutrition and Total Joint Arthroplasty. *J Nat Sci* 2016;2:179-85.
2. Deren ME, Huleatt J, Winkler MF, Rubin LE, Salzler MJ, Behrens SB. Assessment and Treatment of Malnutrition in Orthopaedic Surgery. *JBJS Rev* 2014;2:1-9.
3. Hill R, Paulus S, Dey P, Hurley MA, Carter B. Is undernutrition prognostic of infection complications in children undergoing surgery? A systematic review. *J Hosp Infect* 2016;93:12-21.
4. Vellas B, Villars H, Abellan G, Soto ME, Rolland Y, Guigoz Y, et al. Overview of the MNA® - Its History and Challenges. *J Nut Health Aging* 2006;10:456-65.
5. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1999;20:250-78.
6. Global Guidelines for the Prevention of Surgical Site Infection. Geneva: World Health Organization; 2016.
7. Khanna K, Yi PH, Sing DC, Geiger E, Metz LN. Hypoalbuminemia Is Associated With Septic Revisions After Primary Surgery and Postoperative Infection After Revision Surgery. *Spine (Phila Pa 1976)* 2018;43:454-60.
8. Yuwen P, Chen W, Lv H, Feng C, Li Y, Zhang T, et al. Albumin and surgical site infection risk in orthopaedics: a meta-analysis. *BMC Surg* 2017;17:7.

Table 1. Comparing patients with and without infections with emphasis on nutritional parameters

	Infection		No infection
n = 41	n = 23	p value*	n = 18
Female sex	12 (52%)	0.89	9 (50%)
Median age	74 years	0.49	74.5 years
Immune suppression	7 (30%)	0.57	7 (39%)
- Diabetes mellitus	7 (30%)	0.84	6 (33%)
Elective surgery	2 (9%)	0.22	4 (22%)
Weight (median)	80 kg	0.13	72 kg
- weight loss in the last 3 months (median)	3 kg	0.21	0 kg
Body mass index (median)	28.9 kg/mm ²	0.45	24.3 kg/mm ²
Gastrointestinal disease	3 (13%)	0.85	2 (11%)
Serum albumin (median)	32 g/L	0.11	33 g/L
Serum pre-albumin (median)	170 g/L	0.89	174 g/L
Serum proteins (median)	65 g/L	0.31	63 g/L
Finishes meal*	100%	0.47	100%
MNA questionnaire (median)°	9 points	0.79	9 points

* Percentage of meal finished according to the (auxiliary) nurses

Pearson- χ^2 and Wilcoxon-ranksum-tests, as appropriate

° Mini Nutritional Assessment⁴